

Pathology and Laboratory Medicine

## BLASTOMYCES ANTIGEN

# **Orderable – BLASTOAG**

Turnaround Time: 2 days from receipt at reference laboratory

## **Specimen and Collection Information:**

Please review the MiraVista Diagnostics Laboratory Test Information Index:

#### Blastomyces Quantitative EIA Test

Specimen Type	Minimum Volume	Collection Kit
Urine	0.5 mL	Sterile Container
Serum	1.2 mL	Gold Vacutainer (SST)



Microbiology

Laboratory:

**Requisition:** 

**Ⅲ**×⊓

Method of Analysis:
Quantitative Sandwich
Enzyme Immunoassay
(EIA)

#### **Limitations**

- Result is not intended to be used as the sole means for clinical diagnosis or patient management decisions
- Cross-reactions are seen with histoplasmosis, paracoccidioidomycosis, penicilliosis, less frequently in coccidioidomycosis, rarely in aspergillosis and possibly sporotrichosis



Test Schedule: Referred out weekdays Comments:



### \_\_\_\_\_

Testing restricted to Infectious Diseases (ID) service following a Microbiologist consultation.

#### Referred to:

MiraVista Diagnostics 4705 Decatur Blvd. Indianapolis, Indiana 46241 USA