



Pathology and Laboratory Medicine

Orderable - MACPRLRO

Turnaround Time: 7 days

Specimen:

5 mL Gold top or 6 mL Red top Vacutainer tube Lithium heparin plasma is also acceptable

Collection Information:

2 mL of serum/plasma are requested. Minimum volume of serum/plasma required is 0.4 mL.



Laboratory: Core Lab



Requisition: GENERAL LABORATORY REQUISITION

	•
•	
1	7

Method of Analysis: Immunoenzymatic

Test Schedule: Referred out Monday-

Thursday

Reference Ranges:

Prolactin	
Male:	Less than 25 ug/L
Female:	Less than 18 ug/L

Macroprolactin
Normal: <0.200
Borderline: 0.200 - 0.400
Abnormal: >0.400

Interpretive Comments:

Assay includes Prolactin.



Referred Out Location: In-Common Laboratories

Comments:

To order macroprolactin testing, prolactin testing in our laboratory must be ordered concurrently or have been performed within the last 30 days. Macroprolactin testing should only be ordered if the patient has an elevated prolactin result that is inconsistent with the clinical status. If the prolactin level is not \geq 33 and <470 µg/L, the macroprolactin test is not permitted.

Storage and Shipment:

Store and send frozen.