



Pathology and Laboratory Medicine

PNEUMOCOCCAL ANTIBODY TITRE

# Orderable - PNEUMA(Pre), PNEUMB (post), PNEUM

**Turnaround Time: 6-8 weeks** 

#### **Alternate Name(s):**

Anti-Pneumococcal capsular polysaccharide Ab, IgG, serum PCP

ANP

Pneumococcal Antibody Titre, Pre Vaccine Pneumococcal Antibody Titre, Post Vaccine Pneumococcal Antibody Titre, Single Sample

#### **Specimen:**

Adult	Pediatric
6 mL Red top Vacutainer	0-2 years: 0.5 mL Red top
tube	Microtainer
	2-10 years: 2 mL Red top
	Vacutainer tube

### **Collection Information:**

3 mL serum required. Must indicate "pre" or "post".

Serum from gold top tube not acceptable.

Minimum volume of serum is 0.5 mL

<u>Note:</u> This test is only available at the request of Dr. Ashley Geerlinks, Dr. Kara Robertson, Dr. April Price, Dr. Mark Kuprowski, Dr. Harold Kim, Dr. Rongbo Zhu and Dr. Samira Jeimy.



## Method of Analysis:

**Laboratory:** Core Lab

Requisition:

VaccZyme Anti-PCP IgG enzyme Immunoassay Kit

<u>Biochemistry Immunology</u> Requisition (sickkids.ca)

# \_\_\_

#### **Test Schedule:**

Referred out as required.

#### Referred Out Location:

The Hospital for Sick Children Rapid Response Laboratory- Room 3642 555 University Avenue, Toronto, ON M5G 1X8 Canada

Phone: 416-813-7200 Toll Free: 1-855-381-3212

#### **Reference Ranges:**

Age (years)	Reference Interval (mg/L)
0.5-<1	0.9-93.0
1-<2	0.9-29.2
2-<3	1.4-110.4
3-<4	0.8-262.1
4-<8	9.2-225.9







Pathology and Laboratory Medicine

8-<12	11.0-320.8
12-<18	8.7-502.6
≥18	10.0-191.2

Schauer U et al. Clinical and Diagnostic Laboratory Immunology. 2003. p202-207.

If the patient received Prevnar, anti-pneumococcal levels reported do not represent a response to polysaccharide antigens. This assay tests anti-pneumococcal antibodies, which detect a mix of various serotypes. It, therefore, cannot exclude poor to no response to one or more specific pneumococcal serotypes.

#### **Special Processing:**

Aliquot serum and freeze.

Write the test in the SickKids Clinical Information/Diagnosis box. If you need both a pre and post vaccine value, please indicate that as well and provide both samples.

#### **Storage and Shipment:**

Freeze serum at -20°C. Ship frozen.