

MYCOPLASMA/ UREAPLASMA DETECTION

Orderable - MYPC

Turnaround Time: Up to 17 days

Please review the Public Health Ontario Laboratory Test Information Index:

[Ureaplasma/Mycoplasma Culture](#)

Specimen Type and Collection Information:

Specimen Type	Minimum Volume	Collection Kit
Genital Swabs, including endometrial, placental swabs	1.0 mL of fluid	Viral Transport (VTM) for swabs or fluids.
Neonatal specimens – auger suction, nasopharyngeal/endotracheal aspirates	As much fluid as possible; if <1.0 mL, transfer to a UTM	Sterile Container can also be used for fluids/tissues.
Urine	1.0 mL	
Autopsy tissues/swabs	Available tissue	
Placenta, products of conception	Available Tissue	
Body Fluids, wound exudates/aspirates, joint fluids, needle aspirates	1.0 mL of fluid	



Laboratory:
Microbiology



Requisition:
See electronic order



Method of Analysis:
Culture



Test Schedule:
Referred out weekdays



Referred to:
Public Health Laboratory
1200 Commissioners Rd E
London, Ontario
519-455-9310

Comments:

Unsuitable specimens: Rectal and throat swabs, and seminal fluid are not acceptable specimen sites for Ureaplasma/Mycoplasma culture.


Speciation is not available by routine culture methods.

Antibiotic sensitivity testing is not available for these organisms.

Mycoplasma genitalium: Not detected by culture methods. Molecular testing is available for patients that meet the following criteria

- Symptomatic, in the presence of persistent or recurrent urethritis, cervicitis, or PID
- Negative test results for *Chlamydia trachomatis* and *Neisseria gonorrhoeae*

In specimen comment, include mycoplasma genitalium and confirm test negative for *Chlamydia trachomatis* and *Neisseria gonorrhoeae*



MYCOPLASMA/
UREAPLASMA
DETECTION

Storage and Shipment:

The specimen must be received within 24 hours of collection in the Microbiology Laboratory

Store at 2-8°C if shipped within 48 hours
If stored >48 hours, freeze at -70°C.