

Maintenance Bundle for Maintenance of all Intravascular Devices

Ensure that **patient and health care provider safety standards** are met during this procedure including:

- Risk assessment and appropriate PPE
 - 4 Moments of Hand Hygiene
 - Two patient identification
 - Safe patient handling practices
 - Biomedical waste disposal policies
1. Review insertion date, circumstance and need for continued line use Q shift.
 2. Change lines (arterial, central venous and peripheral) inserted without full precautions (e.g. during emergency situations) within 24-48 hrs (document plan)
 3. Palpate and visually inspect site daily
 4. Ensure catheter securement; change/remove positional lines
 5. Change transparent dressings Q7 days and PRN if soiled, integrity is disrupted, edges are curled or CHG pad feels “boggy” .
 6. Change gauze dressings daily and inspect site
 7. Apply needleless access device to all injection and blood sampling ports (connect pressure tubing directly) to catheter hub.
 8. Back flush sampling ports (into vacuum tube) after blood drawing. Replace needleless access device when soiled or blood remains present
 9. Apply new antiseptic cap to all injection and sampling ports and to the male end of any vascular tubing during temporary disconnection after each access.
 10. Scrub the hub and allow 30 second dry time before accessing ports without antiseptic cap.
 11. Flush lines thoroughly after blood sampling. Flush EACH PICC lumen with 20 ml using turbulent flushing (stop/start technique) after blood sampling or each time a locked device is accessed
 12. Routine tubing changes: a) TPN and insulin Q 24 hrs, b) blood tubing after 2 units (except rapid infuser), c) propofol bottle and tubing Q12 hrs, d) all other sets Q 96 hrs and PRN .
 13. Maintain dedicated line for TPN
 14. Don non-sterile gloves and do not touch insertion site after skin prep for venipuncture and peripheral IV insertion. Venipuncture and peripheral IVs can be source for central line infection.
 15. Maintain aseptic technique for peripheral IVs and document compliance
 16. Document assessment findings in the intravascular section of the 24 Hour Flow sheet Q shift and PRN and update Kardex.
 17. Blood cultures:
 - a) Minimum of 2 sets for any culture event
 - b) Change needleless access cap BEFORE blood culture sampling
 - c) Include discard sample UNLESS IT CONTAINS CITRATE (e.g. dialysis lines)
 - d) If line > 48 hrs, send venipuncture AND line culture(s) and request “CAB” assessment. Draw and order all samples within a 15 minute timeframe and send all bottles in one bag (or bags wrapped together)
 - e) Identify catheter site and type (e.g., R IJ HD) and date of central and arterial catheter insertion (including PICC/HD lines) when ordering cultures

Every member of the team is expected to remind others/stop procedures if any steps are overlooked.