

Prehospital Medical Directives: How Often Are They Being Applied When Indicated: A Retrospective Review

A. Sereda¹, T. Vandeyar¹, M. Davis^{1,2}, A. Dukelow^{1,2}, S. McLeod¹, S. Rodriguez², M. Lewell^{1,2}

¹Division of Emergency Medicine, Faculty of Medicine, The University of Western Ontario, London, Canada

²Southwest Ontario Regional Base Hospital Program, London Health Sciences Centre, London, Canada



Introduction

- Prehospital medical directives are used by EMS personnel to initiate medical treatment before patients arrive in the emergency department
- Appropriate application of these directives can improve patient outcomes by starting treatments at first patient contact
- It is unknown whether prehospital directives are applied to all eligible patients

Objectives

- To determine the proportion of patients transported by EMS who where eligible for medical directive care
- To determine the proportion of eligible patients who received directive based care

Methods

- A retrospective medical record review was performed for patients ≥ 18 years of age brought to the ED via EMS over a 6 month period
- Paramedics' application of a prehospital medical directive was obtained from the Ambulance Call Record (ACR) and classified according to the local 2005 "Primary Care Paramedic Medical Directives
- Patient's chief complaint and vital signs were collected from the ACR using a standardized data extraction tool
- Investigators used this information to determine whether patients met the criteria for the application of the appropriate prehospital medical directive

Results







Figure 1. Proportion of patients requiring pre-hospital medical directives



- Paramedics correctly identified 18 out of 21 (86%) patients who qualified for the administration of a prehospital directive
- No patients were incorrectly identified as requiring a prehospital medical directive




Figure 2. Medical directives utilized

Limitations

Did not capture no service calls in the sample

Conclusions

- Paramedics are identifying patients who meet the criteria for prehospital medical directives with a high level of accuracy
- Paramedics are not applying prehospital medical directives in patients who do not meet the required criteria for those medical directives