

ADRENAL ANTIBODIES, SERUM

Orderable – GENX: then Adrenal Antibodies

Turn Around Time: 10 days

Alternate Name(s):

Anti adrenal antibodies
Steroid Cell Antibodies

Specimen:

5 mL Gold top Vacutainer tube

Collection Information:

Minimum volume of serum required is 1 mL

Grossly hemolyzed, lipemic, or microbially contaminated specimens are unsuitable.

Note: Test approved for Endocrinologists. For all other requests, please page the Biochemist on-call.

Avoid repeated freezing and thawing of samples. Analysis and report include Mitochondrial Antibodies. In the presence of positive Mitochondrial Antibodies (as confirmed on rat kidney cells), positive Adrenal Antibodies cannot be reported and will be noted on report.

Reference Ranges:

Adrenal Antibodies:	Mitochondrial Antibodies:
Negative	Negative

Special Processing:

LTC Staff:
Collect sample, manually label and send to SRA to order



Laboratory:
Core Lab



Requisition:
GENERAL LABORATORY
REQUISITION




Method of Analysis:
Indirect
Immunofluorescence
Assay (IFA)



Test Schedule:
Referred out Monday-
Thursday



Referred Out Location:
[In-Common Laboratories](#)



ADRENAL
ANTIBODIES,
SERUM

Specimen Stability:

Refrigerated: 1 week

Frozen: > 1 week

Additional: Avoid freeze/thaw cycles

Comments:

Note: Test approved for Endocrinologists. For all other requests, please page the Biochemist on-call.