

ANDROSTENEDIONE,
SERUM

Orderable - ANDRO

Turn Around Time: 15 days

Specimen:

Adult	Pediatric
6 mL Red top Vacutainer tube	0-2 years: 2 x 0.5 mL Red top Microtainers 2-10 years: 2 x 0.5 mL Red top Microtainers
Plasma (heparin) is also acceptable	



Laboratory:
Core Lab



Requisition:
GENERAL LABORATORY
REQUISITION



Method of Analysis:
LC-MS/MS



Test Schedule:
Referred out Monday-
Thursday



Referred Out Location:
[In-Common Laboratories](#)

Collection Information:

Minimum volume of serum (or plasma) required is 0.5 mL.

Reference Ranges:

Male	
Age	Range
0 – <14 days:	0.0 – 2.5 nmol/L
14 days – <1 year:	0.1 – 2.1 nmol/L
1 – <6 years	0.1 – 0.6 nmol/L
6 – <10 years:	0.2 – 0.9 nmol/L
10 – <12 years:	0.0 – 2.5 nmol/L
12 – <15 years	0.5 – 2.0 nmol/L
15 – <19 years:	0.9 – 3.6 nmol/L
19 – <31 years:	1.8 – 7.7 nmol/L
31 – <51 years:	1.4 – 6.6 nmol/L
51 – <89 years:	1.8 – 7.7 nmol/L

Female	
Age	Range
0 – <14 days:	0.0 – 2.5 nmol/L
14 days – <1 year:	0.1 – 2.1 nmol/L
1 – <6 years	0.1 – 0.6 nmol/L

ANDROSTENEDIONE,
SERUM

6 – <10 years:	0.2 – 0.9 nmol/L
10 – <12 years:	0.0 – 2.5 nmol/L
12 – <15 years:	0.7 – 6.0 nmol/L
15 – <19 years:	0.5 – 6.5 nmol/L
Follicular Phase:	1.2 – 8.7 nmol/L
Midcycle:	2.1 – 10.0 nmol/L
Luteal Phase:	1.1 – 8.2 nmol/L
Post-menopausal:	0.7 – 2.6 nmol/L

Storage and Shipment:

Store and ship serum (or plasma) frozen.

Specimen Stability:

Refrigerated: 2 days

Frozen: 2 months