

# GLIADIN (DGP) ANTIBODIES

## Orderable - GLIAB

Turn Around Time: 5 days

### Alternate Name(s):

Anti-Deamidated Gliadin Peptide

### Specimen:

6 mL Red top Vacutainer tube

### Collection Information:

Collect blood aseptically in a Vacutainer tube.

### **Note:**

This test is not to be used for Celiac screening unless TTGAb negative and patient IgA deficient - Call the clinical biochemist at 77626 or biochemist on call, Monday to Friday 08:00 to 16:00 for approval before ordering.

### Reference Ranges:

For Deamidated Gliadin Antibody IgA and IgG:	
<20 CU:	Negative
20-30 CU:	Weakly Positive
>30 CU:	Positive

### Interpretive Comments:

The test done for Celiac Disease is Tissue Transglutamase Antibodies.

For screening for diagnosis of Celiac Disease, the patient should not be on a gluten free diet.

False negative results may occur in individuals who are IgA-deficient.

### Storage and Shipment:



**Laboratory:**  
Core Lab



**Requisition:**  
GENERAL LABORATORY  
REQUISITION



**Method of Analysis:**  
CLIA



**Test Schedule:**  
Referred out Monday -  
Thursday



**Referred Out Location:**  
[In-Common Laboratories](#)



GLIADIN  
(DGP)  
ANTIBODIES

Store and ship frozen.